arXiv:math/0206278v1 [math.AP] 26 Jun 2002

On 2D Euler Equations: III. A Line Model

Yanguang (Charles) Li

Abstract. The spectral theorem of the linear 2D Euler operator in Sobolev spaces is presented as a corollary of the spectral theorem in ℓ_2 space in [**1**]. Study on the (dashed) line model introduced in [**2**] is continued. Specifically, invariant manifolds for the line model is established. The corresponding line model for 2D Navier-Stokes equation is also introduced.

1. Introduction

To understand the nature of turbulence, we select 2D Euler equation under periodic boundary condition as our primary example to study. 2D Navier-Stokes equation at high Reynolds number is regarded as a singularly perturbed 2D Euler equation. That is, we are interested in studying the zero viscosity limit problem.

To begin an infinite dimensional dynamical system study, we consider a simple fixed point and study the spectrum of the linear 2D Euler operator in [**1**]. The spectral theorem in ℓ_2 space is proved. As a corollary of the spectral theorem in ℓ_2 space, we will present the spectral theorem in Sobolev spaces in this article. Sobolev spaces are of more interest to us, since we are interested in understanding the invariant manifolds of 2D Euler equation at the fixed point. The main obstacle toward proving the invariant manifold theorem is that the nonlinear term is non-Lipschitzian. In [**2**], a (dashed) line model is introduced to understand the invariant manifold structure of 2D Euler equation. At a special parameter value, the explicit expression of the invariant manifolds of the dashed line model can be calculated. The stable and unstable manifolds are two dimensional ellipsoidal surfaces, and together they form a lip-shape hyperbolic structure. Such structure appears to be robust with repsect to the parameter. In this article, we will prove the existence of invariant manifolds for the line model.

Another more exciting development is the discovery of a Lax pair for 2D Euler equation [**3**]. From the Lax pair, we have obtained a Darboux transformation for the 2D Euler equation [**4**]. In principle, explicit expressions of the hyperbolic structures can be obtained from Darboux transformations [**5**].

1991 *Mathematics Subject Classification.* Primary 76, 35; Secondary 34, 37.

Key words and phrases. Euler equation, invariant manifold, line model, point spectrum, continuous spectrum.

.

©0000 (copyright holder)

Hyperbolic structures are the source of chaos when the system is under perturbations. The corresponding line model for 2D Navier-Stokes equation with temporally periodic forcing is a singular perturbation of the line model for 2D Euler equation. Numerical simulations on these line models are conducted.

The article is organized as follows: In section 2, we present the spectral theorem in Sobolev spaces. In section 3, we present the invariant manifold result on the line model. In section 4, we introduce the corresponding line model for 2D Navier-Stokes equation.

2. The Spectral Theorem of the Linear 2D Euler Operator in Sobolev Spaces

Consider the 2D Euler equation in vorticity form

$$\frac{\partial \Omega}{\partial t} + \{\Psi, \Omega\} = 0 \ , \tag{2.1}$$

where Ω is the vorticity, Ψ is the stream function, $\Omega = \Delta \Psi$, Δ is the 2D Laplacian, and the bracket $\{\ ,\ \}$ is defined as

$$\{f, g\} = (\partial_x f)(\partial_y g) - (\partial_y f)(\partial_x g) \ .$$

Expanding Ω into Fourier series,

$$\Omega = \sum_{k \in \mathbb{Z}^2/\{0\}} \omega_k \ e^{ik\cdot X} \ ,$$

where $\omega_{-k} = \overline{\omega_k}$, $k = (k_1, k_2)^T$, and $X = (x, y)^T$. The 2D Euler equation can be rewritten as

$$\dot{\omega}_k = \sum_{k=p+q} A(p, q) \ \omega_p \omega_q \ , \tag{2.2}$$

where $A(p, q)$ is given by,

$$\begin{aligned} A(p,q) &= \frac{1}{2}[|q|^{-2} - |p|^{-2}](p_1 q_2 - p_2 q_1) \\ &= \frac{1}{2}[|q|^{-2} - |p|^{-2}] \begin{vmatrix} p_1 & q_1 \\ p_2 & q_2 \end{vmatrix} \ , \end{aligned} \tag{2.3}$$

where $|q|^2 = q_1^2 + q_2^2$ for $q = (q_1, q_2)^T$, similarly for p. Denote $\{\omega_k\}_{k \in \mathbb{Z}^2/\{0\}}$ by ω. We consider the simple fixed point ω^*:

$$\omega_p^* = \Gamma, \quad \omega_k^* = 0, \text{ if } k \neq p \text{ or } -p, \tag{2.4}$$

of the 2D Euler equation (2.2), where Γ is an arbitrary complex constant. The *linearized two-dimensional Euler equation* at ω^* is given by,

$$\dot{\omega}_k = A(p, k-p) \ \Gamma \ \omega_{k-p} + A(-p, k+p) \ \bar{\Gamma} \ \omega_{k+p} \ . \tag{2.5}$$

Definition 2.1 (Classes). For any $\hat{k} \in \mathbb{Z}^2/\{0\}$, we define the class $\Sigma_{\hat{k}}$ to be the subset of $\mathbb{Z}^2/\{0\}$:

$$\Sigma_{\hat{k}} = \left\{ \hat{k} + np \in \mathbb{Z}^2/\{0\} \ \middle| \ n \in \mathbb{Z}, \quad p \text{ is specified in (2.4)} \right\}.$$

See Fig.1 for an illustration of the classes. According to the classification defined in Definition 2.1, the linearized two-dimensional Euler equation (2.5) decouples into infinitely many *invariant subsystems*:

$$\begin{aligned} \dot{\omega}_{\hat{k}+np} &= A(p, \hat{k}+(n-1)p)\ \Gamma\ \omega_{\hat{k}+(n-1)p} \\ &\quad + A(-p, \hat{k}+(n+1)p)\ \bar{\Gamma}\ \omega_{\hat{k}+(n+1)p}\ . \end{aligned} \tag{2.6}$$

Let $\mathcal{L}_{\hat{k}}$ be the linear operator defined by the right hand side of (2.6), and H^s be the Sobolev space where $s \geq 0$ is an integer and $H^0 = \ell_2$.



FIGURE 1. An illustration of the classes $\Sigma_{\hat{k}}$ and the disk $\bar{D}_{|p|}$.

THEOREM 2.2. *The eigenvalues of the linear operator* $\mathcal{L}_{\hat{k}}$ *in* H^s *are of four types: real pairs* $(c, -c)$*, purely imaginary pairs* $(id, -id)$*, quadruples* $(\pm c \pm id)$*, and zero eigenvalues.*

Proof. The same proof as in [**1**] works here. QED
The eigenvalues can be computed through continued fractions [**1**].

DEFINITION 2.3 (The Disk). The disk of radius $|p|$ in $\mathbb{Z}^2/\{0\}$, denoted by $\bar{D}_{|p|}$, is defined as

$$\bar{D}_{|p|} = \left\{ k \in \mathbb{Z}^2/\{0\} \ \middle|\ |k| \leq |p| \right\} .$$

See Fig.1 for an illustration.

THEOREM 2.4 (The Spectral Theorem). *We have the following claims on the spectrum of the linear operator* $\mathcal{L}_{\hat{k}}$*:*

1. *If* $\Sigma_{\hat{k}} \cap \bar{D}_{|p|} = \emptyset$, *then the entire* H^s *spectrum of the linear operator* $\mathcal{L}_{\hat{k}}$ *is its continuous spectrum. See Figure 2, where* $b = -\frac{1}{2}|\Gamma||p|^{-2} \begin{vmatrix} p_1 & \hat{k}_1 \\ p_2 & \hat{k}_2 \end{vmatrix}$. *That is, both the residual and the point spectra of* $\mathcal{L}_{\hat{k}}$ *are empty.*
2. *If* $\Sigma_{\hat{k}} \cap \bar{D}_{|p|} \neq \emptyset$, *then the entire essential* H^s *spectrum of the linear operator* $\mathcal{L}_{\hat{k}}$ *is its continuous spectrum. That is, the residual spectrum of* $\mathcal{L}_{\hat{k}}$ *is empty. The point spectrum of* $\mathcal{L}_{\hat{k}}$ *is symmetric with respect to both real and imaginary axes. See Figure 3.*

Proof. The same proof as in [**1**] can be carried through here for H^s, with the following minor modifications:

1. In the proof of Theorem VI.1 on page 747 of [**1**], simply replace the ℓ_2-norm by H^s-norm.
2. In the proof of Theorem VI.3 on page 750 of [**1**], the inner product $\langle\ ,\ \rangle$ should still be an ℓ_2 inner product.
3. In the proof of Theorem VI.4 on page 751 of [**1**], ℓ_1, ℓ_2, and ℓ_∞ should be replaced by the Sobolev spaces $W^{s,1}$, $W^{s,2}$ ($= H^s$ in our notation), and $W^{s,\infty}$. In the expression (VI.60) of f_n on page 753, one has
$$\begin{aligned} n^s f_n \quad = \quad & \frac{2\left[1 - w_*^{-4}\right]}{W_0} \sum_{j=0}^{n-1} \left[(n-j)^s w_*^{n-j} + (n-j)^s(-w_*)^{n-j}\right] \\ & \left(\frac{n^s}{(n-j)^s(j+2)^s}\right)(j+2)^s y_{j+2} - \frac{2\left[1 - w_*^{4}\right]}{W_0} \sum_{j=n}^{\infty} \\ & \left[w_*^{j-n} + (-w_*)^{j-n}\right] \left(\frac{n^s}{(j+2)^s}\right)(j+2)^s y_{j+2} \, , \end{aligned}$$
where both $\left(\frac{n^s}{(n-j)^s(j+2)^s}\right)$ and $\left(\frac{n^s}{(j+2)^s}\right)$ are bounded in n and j, and the Riesz convexity theorem can be applied.

The proof of the theorem is completed. QED

3. Invariant Manifolds of the Line Model

To simplify our study, we study only the case when ω_k is real, $\forall k \in \mathbb{Z}^2/\{0\}$, i.e. we only study the cosine transform of the vorticity,
$$\Omega = \sum_{k \in \mathbb{Z}^2/\{0\}} \omega_k \cos(k \cdot X) \ ,$$
and the 2D Euler equation (2.1) preserves the cosine transform. To further simplify our study, we will study a concrete line model based upon the fixed point (2.4) with the mode $p = (1,1)^T$ and parametrized by Γ. When $\Gamma \neq 0$, the fixed point has 4 eigenvalues which form a quadruple. These four eigenvalues appear in the invariant linear subsystem labeled by $\hat{k} = (-3,-2)^T$. We computed the eigenvalues through continued fractions, one of them is [**1**]:
$$\tilde{\lambda} = 2\lambda/|\Gamma| = 0.24822302478255 \ + \ i \ 0.35172076526520 \, . \tag{3.1}$$
See Figure 4 for an illustration. The essential spectrum (= continuous spectrum) of $\mathcal{L}_{\hat{k}}$ with $\hat{k} = (-3,-2)^T$ is the segment on the imaginary axis shown in Figure 4, where $b = -\frac{1}{4}\Gamma$. The essential spectrum (= continuous spectrum) of the linear



FIGURE 2. The spectrum of $\mathcal{L}_{\hat{k}}$ in case (1).



FIGURE 3. The spectrum of $\mathcal{L}_{\hat{k}}$ in case (2).

2D Euler operator at this fixed point is the entire imaginary axis. The line model



FIGURE 4. The spectrum of $\mathcal{L}_{\hat{k}}$ with $\hat{k} = (-3,-2)^T$, when $p = (1,1)^T$.



FIGURE 5. The collocation of the modes in the line model.

is a Galerkin truncation with the modes on the line $\{\hat{k} + np, \ n \in \mathbb{Z}\}$ and p, where

$\hat{k} = (-3,-2)^T$ and $p = (1,1)^T$. See Figure 5 for an illustration of the modes in this model, which has the line nature leading to the name of the model. The line model is designed to model the hyperbolic structure in the neighborhood of the fixed point. For simplicity of presentation, we use the abbreviated notations,

$$\omega_n = \omega_{\hat{k}+np}\,, \quad A_n = A(p, \hat{k}+np)\,, \quad A_{m,n} = A(\hat{k}+mp, \hat{k}+np)\,.$$

The *line model* is,

$$\begin{aligned} \dot{\omega}_n &= A_{n-1}\omega_p\omega_{n-1} - A_{n+1}\omega_p\omega_{n+1}\ , \\ \dot{\omega}_p &= -\sum_{n\in Z} A_{n-1,n}\omega_{n-1}\omega_n\,. \end{aligned} \tag{3.2}$$

We also use ω^* to denote the fixed point of the line model $\{\omega_p = \Gamma;\ \omega_n = 0,\ \forall n \in \mathbb{Z}\}$. The linearized line model at the fixed point ω^* is the same with the invariant subsystem (2.6) with $\hat{k} = (-3,-2)^T$ and $p = (1,1)^T$. In the neighborhood of the fixed point ω^*, the line model can be rewritten as

$$\dot{\omega} = L\omega + Q(\omega)\ , \tag{3.3}$$

where

$$\begin{aligned} &\omega = (\omega_p, \quad \omega_n \quad (n \in \mathbb{Z}))\ , \\ &[L\omega]_n = A_{n-1}\Gamma\omega_{n-1} - A_{n+1}\Gamma\omega_{n+1}\ , \\ &[L\omega]_p = 0\ , \\ &[Q(\omega)]_n = A_{n-1}\omega_p\omega_{n-1} - A_{n+1}\omega_p\omega_{n+1}\ , \\ &[Q(\omega)]_p = -\sum_{n\in\mathbb{Z}} A_{n-1,n}\omega_{n-1}\omega_n\ . \end{aligned}$$

The spectrum of L is given in Figure 4. We have

Theorem 3.1 (Invariant Manifold Theorem). *The line of fixed points ω^* parametrized by Γ of the line model has codimension 2 smooth center-unstable and center-stable manifolds, and codimension 4 smooth center manifold, in H^s $(s \geq 1)$.*

Proof. The spectrum of L in H^s $(s \geq 1)$ is given in Figure 4. $Q(\omega)$ is quadratic in ω for H^s $(s \geq 1)$ is a Banach algebra. In a small neighborhood of ω^*, the existence of smooth center-unstable, center-stable, and center manifolds follows from standard arguments. QED

4. A Line Model for 2D Navier-Stokes Equation

We consider the 2D Navier-Stokes equation with temporally periodic forcing, as a singular perturbation of the 2D Euler equation (2.1),

$$\frac{\partial\Omega}{\partial t} + \{\Psi, \Omega\} = \epsilon[\Delta\Omega + f(t,x,y)]\ , \tag{4.1}$$

where $\epsilon = 1/\mathrm{Re}$ is the inverse of Reynolds number, and $f(t,x,y)$ is periodic in t, periodic in x and y of period 2π, and of spatial mean 0. The corresponding line model is

$$\begin{aligned}\dot{\omega}_n &= A_{n-1}\omega_p\omega_{n-1} - A_{n+1}\omega_p\omega_{n+1} + \epsilon[-\kappa_n^2\omega_n + f_n(t)] \ , \\ \dot{\omega}_p &= -\sum_{n\in Z} A_{n-1,n}\omega_{n-1}\omega_n + \epsilon[-\kappa_p^2\omega_p + f_p(t)] \, ,\end{aligned} \tag{4.2}$$

where f_n and f_p are periodic functions of t, and

$$\kappa_n = |\hat{k} + np| \ , \quad \kappa_p = |p| \ , \quad \hat{k} = (-3,-2)^T \ , \quad p = (1,1)^T \ .$$

References

[1] Y. Li, On 2D Euler equations. I. On the energy-Casimir stabilities and the spectra for a linearized two dimensional Euler equation, *Journal of Mathematical Physics* **41, no.2** (2000), 728.

[2] Y. Li, On 2D Euler equations: part II. Lax pairs and homoclinic strucutres, *Submitted* (2001).

[3] Y. Li, A Lax pair for the 2D Euler equation, *Journal of Mathematical Physics* **42, no.8** (2001), 3552.

[4] Y. Li and A. Yurov, Lax pairs and Darboux transformations for Euler equations, **Preprint, available at: http://xxx.lanl.gov/abs/math.AP/0101214, or http://www.math.missouri.edu/~cli** (2000).

[5] Y. Li, Bäcklund-Darboux transformations and Melnikov analysis for Davey-Stewartson II equations, *Journal of Nonlinear Sciences* **10, no.1** (2000), 103.

Department of Mathematics, University of Missouri, Columbia, MO 65211
E-mail address: `cli@math.missouri.edu`